

Path Planning 2

Input file: **standard input**
Output file: **standard output**
Time limit: 3 seconds
Memory limit: 1024 megabytes

There is a grid with n rows and m columns. Each cell of the grid has an integer in it, where $a_{i,j}$ indicates the integer in the cell located at the i -th row and the j -th column.

Let (i, j) be the cell located at the i -th row and the j -th column. You now start from $(1, 1)$ and need to reach (n, m) . When you are in cell (i, j) , you can either move to its right cell $(i, j + 1)$ if $j < m$ or move to its bottom cell $(i + 1, j)$ if $i < n$.

Let \mathbb{S} be the set consisting of integers in each cell on your path, including $a_{1,1}$ and $a_{n,m}$. Let $\text{mex}(\mathbb{S})$ be the smallest non-negative integer which does not belong to \mathbb{S} . Find a path to minimize $\text{mex}(\mathbb{S})$ and calculate this minimum possible value.

Input

There are multiple test cases. The first line of the input contains an integer T ($1 \leq T \leq 10^4$) indicating the number of test cases. For each test case:

The first line contains two integers n and m ($1 \leq n, m \leq 10^6$, $1 \leq n \times m \leq 10^6$) indicating the number of rows and columns of the grid.

For the following n lines, the i -th line contains m integers $a_{i,1}, a_{i,2}, \dots, a_{i,m}$ ($0 \leq a_{i,j} \leq 10^9$) where $a_{i,j}$ indicates the integer in cell (i, j) .

It's guaranteed that the sum of $n \times m$ of all test cases will not exceed 10^6 .

Output

For each test case output one line containing one integer indicating the minimum possible value of $\text{mex}(\mathbb{S})$.

Example

standard input	standard output
2	1
2 3	3
2 0 1	
0 3 4	
1 5	
100 0 2 0 1	

Note

For the first sample test case there are 3 possible paths.

- The first path is $(1, 1) \rightarrow (1, 2) \rightarrow (1, 3) \rightarrow (2, 3)$. $\mathbb{S} = \{2, 0, 1, 4\}$ so $\text{mex}(\mathbb{S}) = 3$.
- The second path is $(1, 1) \rightarrow (1, 2) \rightarrow (2, 2) \rightarrow (2, 3)$. $\mathbb{S} = \{2, 0, 3, 4\}$ so $\text{mex}(\mathbb{S}) = 1$.
- The third path is $(1, 1) \rightarrow (2, 1) \rightarrow (2, 2) \rightarrow (2, 3)$. $\mathbb{S} = \{2, 0, 3, 4\}$ so $\text{mex}(\mathbb{S}) = 1$.

So the answer is $\min(3, 1, 1) = 1$.

For the second sample test case there is only 1 possible path, which is $(1, 1) \rightarrow (1, 2) \rightarrow (1, 3) \rightarrow (1, 4) \rightarrow (1, 5)$. $\mathbb{S} = \{100, 0, 2, 1\}$ so $\text{mex}(\mathbb{S}) = 3$.