

## Problem J: Captain Obvious and the Rabbit-Man

*“It’s you, Captain Obvious!” – cried the evil Rabbit-Man – “you came here to foil my evil plans!”*

*“Yes, it’s me.” – said Captain Obvious.*

*“But... how did you know that I would be here, on 625 Sunflower Street?! Did you crack my evil code?”*

*“I did. Three days ago, you robbed a bank on 5 Sunflower Street, the next day you blew up 25 Sunflower Street, and yesterday you left quite a mess under number 125. These are all powers of 5. And last year you pulled a similar stunt with powers of 13. You seem to have a knack for Fibonacci numbers, Rabbit-Man.”*

*“That’s not over! I will learn... **arithmetics!**” – Rabbit-Man screamed as he was dragged into custody – “You will **never** know what to expect... Owwww! Not my ears, you morons!”*

*“Maybe, but right now you are being arrested.” – Captain added proudly.*

Unfortunately, Rabbit-Man has now indeed learned some more advanced arithmetics. To understand it, let us define the sequence  $F_n$  (being not completely unlike the Fibonacci sequence):

$$F_1 = 1,$$

$$F_2 = 2,$$

$$F_n = F_{n-1} + F_{n-2} \text{ for } n \geq 3.$$

Rabbit-Man has combined all his previous evil ideas into one master plan. On the  $i$ -th day, he does a malicious act on the spot number  $p(i)$ , defined as follows:

$$p(i) = a_1 \cdot F_1^i + a_2 \cdot F_2^i + \dots + a_k \cdot F_k^i.$$

The number  $k$  and the integer coefficients  $a_1, \dots, a_k$  are fixed. Captain Obvious learned  $k$ , but does not know the coefficients. Given  $p(1), p(2), \dots, p(k)$ , help him to determine  $p(k+1)$ . To avoid overwhelmingly large numbers, do all the calculations modulo a fixed prime number  $M$ . You may assume that  $F_1, F_2, \dots, F_n$  are pairwise distinct modulo  $M$ . You may also assume that there always exists a unique solution for the given input.

### Input

The first line of input contains the number of test cases  $T$ . The descriptions of the test cases follow:

The first line of each test case contains two integers  $k$  and  $M$ ,  $1 \leq k \leq 4000$ ,  $3 \leq M \leq 10^9$ . The second line contains  $k$  space-separated integers – the values of  $p(1), p(2), \dots, p(k)$  modulo  $M$ .

### Output

Print the answers to the test cases in the order in which they appear in the input. For each test case print a single line containing one integer: the value of  $p(k+1)$  modulo  $M$ .

## Example

For an example input	the correct answer is:
2	30
4 619	83
5 25 125 6	
3 101	
5 11 29	

**Explanation:** the first sequence is simply  $5^i \bmod 619$ , therefore the next element is  $5^5 \bmod 619 = 30$ . The second sequence is  $2 \cdot 1^i + 3^i \bmod 101$ .