

Problem B. Well Off

Input file: *standard input*
 Output file: *standard output*
 Time limit: 2 seconds
 Memory limit: 128 mebibytes

Thanks to his hard work, John, formerly Johnny, has become a member of the respectable Well Off Club. The club has n members, numbered from 1 to n ; the i -th club member has a fortune worth x_i . These can be arbitrary numbers, including negative ones. The reason for this is that the club membership is for life, i.e., neither impoverishment nor even bankruptcy results in an ousting.

The club members take pleasure in comparing their asset values, but they shun telling the number directly; it is their custom to engage in a conversation that lets them establish that their fortunes x_i and x_j satisfy one of the inequalities $x_i + x_j > 0$, $x_i - x_j > 0$, $-x_i + x_j > 0$, or $-x_i - x_j > 0$.

Johnny (over-)heard many such inequalities today at the club, but he suspects that some of the members are lying. Help him determine if all the inequalities could possibly be true, i.e., if they can be simultaneously satisfied by some set of fortunes. Write a program that: reads the inequalities heard by Johnny, determines if they are satisfiable, and prints the answer to the standard output.

Input

In the first line of input there are two integers, n and m ($1 \leq n \leq 100\,000$, $1 \leq m \leq 500\,000$), separated by a single space. These specify the number of club members and the number of inequalities relating their fortunes, respectively.

Each of the m lines that follow describes one inequality, encoded by a + or - sign, an integer i ($1 \leq i \leq n$), another + or - sign, and an integer j ($1 \leq j \leq n$), separated by single spaces. These correspond to the inequality $\pm x_i \pm x_j > 0$ (with the signs as preceding i and j in the description). It might happen that $i = j$.

Output

The first and only line of the output should contain a single word: “TAK” if the system of inequalities is satisfiable, and “NIE” otherwise. (Naturally, these are the Polish words for *yes* and *no*.)

Examples

standard input	standard output
3 3 + 1 - 2 - 3 + 1 + 2 - 3	TAK
3 3 + 1 - 2 + 3 - 1 + 2 - 3	NIE

Note

In Sample 1, system of inequalities is satisfiable, e.g. by $x_1 = 3$, $x_2 = 2$, $x_3 = 1$.

In Sample 2, system of inequalities is not satisfiable.