


Delay Time

Input file: standard input
Output file: standard output
Time limit: 1 second
Memory limit: 512 megabytes

Petr and Egor are measuring the gravitational acceleration g on their physics lessons using a special device. An electromagnetic trap holding a metal ball is installed on an arbitrary positive height chosen by the user. As soon as the button is pressed, the timer starts, the electromagnet turns off and the released ball falls on a table. The timer stops immediately when the ball touches the table.



Petr has set the height to h_1 meters and got t_1 seconds on the timer as a result of the experiment. In his turn Egor **changed** the height to h_2 meters and got t_2 seconds shown on the timer. While calculating the value of g they surprisingly got significantly different results, and asked a teacher to clarify the nature of this inconsistency.

As it often happens with this sort of “mystery”, the solution was simple: there was a constant delay of d seconds between the start of the timer and the moment the ball actually starts to fall, caused by the time needed for the electromagnet to lose its grip on the ball. They quickly managed to determine this value from the given information and calculate the correct g . Can you manage to do this in five hours?

Assume that the gravitational acceleration and the delay time are the same in both experiments. Air friction and other effects are considered negligible for the purpose of this experiment.

When the ball with zero initial speed starts to fall affected only by the gravitational acceleration g , the distance it travels in t seconds is given by a well-known formula:

$$\rho(t) = \frac{g \cdot t^2}{2}$$

Input

The first line of the input contains real values h_1 and t_1 — the height selected by Petr, in meters, and the time value he has measured, in seconds ($0 < h_1, t_1 \leq 10$).

The second line contains real values h_2 and t_2 — the values of height in meters and time in seconds for Egor’s experiment ($0 < h_2, t_2 \leq 10, h_1 \neq h_2$).

All values in the input are given with no more than three digits after the decimal point.

It is guaranteed that input data is consistent, that means that **positive** real g and d were used to generate the input.

Output

Output the value of electromagnet delay d in seconds. Your answer will be considered correct if its absolute or relative error doesn't exceed 10^{-4} .

Formally, if your answer is a and the jury's answer is b , the checker will accept your answer if $\frac{|a-b|}{\max(1,b)} \leq 10^{-4}$.

Examples

standard input	standard output
1.495 2.009 7.674 2.708	1.456709
5 1.25 1.25 0.75	0.250000

Note

In the first sample the gravitational acceleration g is approximately equal to 9.80246, while in the second sample it is 10.

Note that g in the testset can vary arbitrarily and is not guaranteed to be related to the gravitational acceleration on the surface of any known planet.