

# Sum of Log

Input file:            **standard input**  
Output file:           **standard output**  
Time limit:            1 second  
Memory limit:         1024 megabytes

Given two non-negative integers  $X$  and  $Y$ , determine the value of

$$\sum_{i=0}^X \sum_{j=[i=0]}^Y [i \& j = 0] [\log_2(i + j) + 1]$$

modulo  $10^9 + 7$  where

- $\&$  denotes bitwise AND;
- $[A]$  equals 1 if  $A$  is true, otherwise 0;
- $\lfloor x \rfloor$  equals the maximum integer whose value is no more than  $x$ .

## Input

The first line contains one integer  $T$  ( $1 \leq T \leq 10^5$ ) denoting the number of test cases.

Each of the following  $T$  lines contains two integers  $X, Y$  ( $0 \leq X, Y \leq 10^9$ ) indicating a test case.

## Output

For each test case, print one line containing one integer, the answer to the test case.

## Example

standard input	standard output
3	14
3 3	814
19 26	278
8 17	

## Note

For the first test case:

- Two  $(i, j)$  pairs increase the sum by 1:  $(0, 1), (1, 0)$
- Six  $(i, j)$  pairs increase the sum by 2:  $(0, 2), (0, 3), (1, 2), (2, 0), (2, 1), (3, 0)$

So the answer is  $1 \times 2 + 2 \times 6 = 14$ .